Proposed Maximum Residue Limit

PMRL2009-12

Azoxystrobin

(publié aussi en français)

8 December 2009

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications Pest Management Regulatory Agency Health Canada 2720 Riverside Drive A.L. 6605C Ottawa, Ontario K1A 0K9

Internet: pmra.publications@hc-sc.gc.ca healthcanada.gc.ca/pmra Facsimile: 613-736-3758 Information Service: 1-800-267-6315 or 613-736-3799 pmra.infoserv@hc-sc.gc.ca



HC Pub: 091179

ISBN: 978-1-100-14357-6 (978-1-100-14358-3))

Catalogue number: H113-24/2009-12E (H113-24/2009-12E-PDF)

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health Canada, 2009

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.

Under the authority of the *Pest Control Products Act*, Health Canada's Pest Management Regulatory Agency (PMRA) has concluded that the addition of new uses on cereal grains (Crop Group 15; see Appendix I) to the product labels of Quadris Flowable Fungicide and Quilt Fungicide, containing technical grade azoxystrobin, is acceptable. The specific uses approved in Canada are detailed on the labels of Quadris Flowable Fungicide and Quilt Fungicide, *Pest Control Products Act* Registration Numbers 26153 and 28328, respectively.

The evaluation of these azoxystrobin applications indicated that the end-use products have merit and value and that the human health and environmental risks associated with the new uses are acceptable. Details regarding the registration can be found in the corresponding Evaluation Report that is available in the Pesticides and Pest Management section of Health Canada's website, under Public Registry, Pesticide Product Information Database.

Before registering a pesticide for food use in Canada, the PMRA must determine the quantity of residues that are likely to remain in or on the food when the pesticide is used according to label directions and that such residues will not be a concern to human health. This quantity is then legally established as a maximum residue limit (MRL). An MRL applies to the identified raw agricultural food commodity as well as to any processed food product that contains it, except where separate MRLs are specified for the raw agricultural commodity and a processed product made from it.

In addition, the PMRA is consulting on proposed MRLs for azoxystrobin on commodities recently registered in Canada for which MRL consultations have not yet been conducted. These MRLs are included in Table 1 and are supported by the summary residue data captured in Appendix II of this PMRL.

Consultation on the proposed MRLs for azoxystrobin is being conducted via this document (see Next Steps, the last section of this document).

To comply with Canada's international trade obligations, consultation on the proposed MRLs is also being conducted internationally by notifying the World Trade Organization, as coordinated by the Standards Council of Canada.

The relevant report can be accessed by selecting the Applications/Amendment/Historical tab and opening the Evaluation Report found under either Application Number 2006-3665 (Quadris) or 2006-3668 (Quilt).

The proposed MRLs for azoxystrobin in Canada in or on food, to be added to those MRLs already legally established, are as follows.

Table 1 Proposed Maximum Residue Limits for Azoxystrobin

| Common Name | Residue Definition | MRL (ppm) | Food Commodity | | |
|--------------|---|-------------------------|---|--|--|
| Azoxystrobin | (αE)-methyl 2-[[6- 2-yanophenoxy)-4- pyrimidinyl]oxy]-α- (methoxymethylene) benzeneacetate, including the isomer (Z)-methyl 2-[[6- (2-cyanophenoxy)-4- pyrimidinyl]oxy]-α- (methoxymethylene) benzeneacetate | 0.2* 0.03** 0.02* | Succulent shelled English peas, succulent shelled garden peas, succulent shelled green peas, succulent shelled pigeon peas Barley, buckwheat, oats, pearl millet, popcorn grain, proso millet, rye, sorghum, teosinte, triticale, wheat, wild rice Edible-podded jackbeans, edible-podded moth beans, edible-podded runner beans, edible-podded snap beans, edible-podded soybeans, edible-podded swordbeans, edible-podded wax beans, edible-podded swordbeans, edible-podded wax beans, edible-podded yardlong beans, succulent shelled blackeyed peas, succulent shelled broad beans, succulent shelled cowpeas, succulent shelled lima beans, succulent shelled southern peas | | |

The MRLs proposed for these previously registered commodities are supported by corresponding residue trial data attached under Appendix II.

MRLs for azoxystrobin are already established on field corn (0.03 ppm), rice (4.0 ppm) and sweet corn kernels plus cob with husks removed (0.03 ppm). This PMRL proposes to extend the 0.03 ppm MRL to the remaining cereal grain commodities.

A complete list of all MRLs established in Canada can be found on the Maximum Residue Limits for Pesticides webpage in the Pesticides and Pest Management section of Health Canada's website.

International Situation and Trade Implications

MRLs may vary from one country to another for a number of reasons, including differences in pesticide use patterns and the locations of the field crop trials used to generate residue chemistry data. As per Table 2, the proposed MRLs in Canada differ from the corresponding tolerances established in the United States (tolerances are listed in the Electronic Code of Federal Regulations by pesticide). Currently, Codex² MRLs have not been established for azoxystrobin on any commodity. A listing of all established Codex MRLs is available on the Pesticide Residues in Food website.

Codex is an international organization under the auspices of the United Nations that develops international food standards, including MRLs.

Table 2 Comparison of Canadian MRLs, American Tolerances and Codex MRLs

| Food Commodity | Canadian MRL (ppm) | American Tolerance (ppm) | Codex MRL (ppm) | |
|--|--------------------|-----------------------------|--------------------|--|
| Succulent shelled pea and bean (Crop subgroup 6B), pea commodities | 0.2 | 0.5 | No MRL established | |
| Barley | 0.03 | 3.0 | * | |
| Barley bran | 0.03* 6.0 | | | |
| Buckwheat | 0.03 | No tolerance established | | |
| Oats | 0.03 | No tolerance established | | |
| Pearl millet | 0.03 | No tolerance established | | |
| Popcorn grain | 0.03 | 0.05 | | |
| Proso millet | 0.03 | No tolerance established | | |
| Rye | 0.03 | No tolerance established | | |
| Sorghum | 0.03 | 11 | | |
| Teosinte | 0.03 | No tolerance established | | |
| Triticale | 0.03 | 0.1 | | |
| Wheat | 0.03 | 0.1 | | |
| Wheat bran | 0.03* | 0.2 | | |
| Wild rice | 0.03 | 5.0 | | |
| Edible-podded legume vegetables (Crop subgroup 6A), bean commodities | 0.02 | 3.0 | | |
| Succulent shelled pea and bean (Crop subgroup 6B), bean commodities | 0.02 | 0.5 | | |

Covered by the MRL proposed for the raw agricultural commodities barley and wheat.

Next Steps

The PMRA invites the public to submit written comments on the proposed MRLs for azoxystrobin up to 75 days from the date of publication of this document. Please forward your comments to Publications (see the contact information on the cover page of this document). The PMRA will consider all comments received before making a final decision on the proposed MRLs for azoxystrobin and posting a corresponding Established Maximum Residue Limit in the Pesticides and Pest Management section of Health Canada's website.

Proposed Maximum Residue Limit - PMRL2009-12 Page 4

Appendix I Crop Groups: Numbers and Definitions

| Crop Group Number | Crop Group Name | Food Commodities Included in the Crop Group | | | | |
|----------------------|-----------------|--|--|--|--|--|
| 15 | Cereal grains | Barley Buckwheat Field corn* Oats Pearl millet Popcorn grain Proso millet Rice* Rye Sorghum Sweet corn kernels plus cob with husks removed* Teosinte Triticale Wheat Wild rice | | | | |

MRLs for these cereal grains are already established and they are therefore excluded from this action.

| Α | | | |
|---|--|--|--|
| | | | |

Appendix II Summary of Field Trial Data

Table 1 Summary of Field Trial Data Used to Establish Maximum Residue Limits (MRLs) for Bean Commodities in the Edible-podded Legume Vegetables (6A) and Succulent Shelled Pea and Bean (6B) Subgroups.

| Commodity | Application Method/Total Application Target Rate (g a.i./ha) | PHI (days) | Residues ¹ (ppm) | | Recommended MRL for |
|------------------|---|---------------|-----------------------------|-------|--|
| | | | Min | Max | Edible-podded and Succulent Shelled Bear Commodities |
| Bean with pod | Foliar spray / 225–250 | 13-48 | <0.02 | <0.02 | 0.02 ppm |
| Bean without pod | Foliar spray / 225–250 | 13–15 | <0.02 | <0.02 | 0.02 ppm |

Combined residues of azoxystrobin and the z-isomer.

Table 2 Summary of Field Trial Data Used to Establish Maximum Residue Limits (MRLs) for Pea Commodities in the Succulent Shelled Pea and Bean (6B) Subgroup.

| Commodity | Application | PHI | Residues ¹ (ppm) | | Recommended MRL for |
|-----------------------|--|--------|-----------------------------|--------|--------------------------------------|
| | Method/Total Application Target Rate (g a.i./ha) | (days) | Min | in May | Succulent Shelled Pea Commodities |
| Succulent shelled pea | 1818–1863 | 0 | 0.04 | 0.18 | 0.2 ppm |

Combined residues of azoxystrobin and the z-isomer.

NOTE:

g - grams

a.i. - active ingredient

ha - hectare

PHI - preharvest interval

